

Electroweak baryon number violation and a “topological condensate”

Steven Bass / Innsbruck

Baryon asymmetry in the Universe:

Needs baryon number non-conservation

→ Non-perturbative “Sphaleron” processes in the electroweak vacuum as candidate mechanism

Consistency of local Ward identities and non-local topological structure

→ formation of a “topological condensate” in the early Universe

QCD version of this physics provides an elegant “solution” to the proton spin puzzle: quarks contribute just 30% of the proton’s spin

Venice, August 29 2007

Baryon asymmetry in the Universe

- **What is observed ?**

$$\gg n_B / n_{\text{gamma}} = 10^{-10}$$

- **Sakharov's conditions (1966)**

- Baryon number non-conservation
 - (non-perturbative e-weak processes or „new physics“)
- C and CP violation
 - Need different rate of reactions with particles and antiparticles
- Deviations from thermal equilibrium
 - Otherwise, if the initial baryon number were zero at the start of Universe it would stay zero

- **Here: Focus on sphaleron induced electroweak baryon number violation**

A Key Issue: What is baryon number ?

- Definition of baryon number in e-weak theory is subtle because of the axial anomaly

$$\begin{aligned} J_\mu &= \bar{\Psi} \gamma_\mu \Psi \\ &= \bar{\Psi} \gamma_\mu \frac{1}{2}(1 - \gamma_5) \Psi + \bar{\Psi} \gamma_\mu \frac{1}{2}(1 + \gamma_5) \Psi. \end{aligned}$$

- SU(2) gauge bosons couple only to left handed quarks \rightarrow axial anomaly is important!

$$\partial^\mu J_\mu = n_f (-\partial^\mu K_\mu + \partial^\mu k_\mu),$$

$$\partial^\mu K_\mu = \frac{g^2}{32\pi^2} W_{\mu\nu} \tilde{W}^{\mu\nu}$$

- **Suggests choice of currents to define baryon number:**
(1) the gauge invariant renormalized current
OR
(2) (gauge invariant observables associated with) the conserved (but gauge dependent) current

$$J_\mu^{\text{con}} = J_\mu - n_f (-K_\mu + k_\mu)$$

$$\partial^\mu J_\mu^{\text{con}} = 0$$

Anomalous commutators

- Consider the charges

$$Y(t) = \int d^3z J_0(z), \quad B = \int d^3z J_0^{\text{con}}(z).$$

- Gauge invariant baryon number B is defined through the commutators

$$[B, \mathcal{O}]_- = B\mathcal{O}$$

despite gauge dependence of the operator

- B charge is renormalization scale invariant (as baryon number should be!) whereas Y is not. Also, the time derivative of the spatial components of the W boson field has zero B charge and non-zero Y charge

$$[B, \partial_0 A_i]_- = 0$$

and


$$\begin{aligned} \lim_{t' \rightarrow t} [Y(t'), \partial_0 A_i(\vec{x}, t)]_- \\ = \frac{in_f g^2}{4\pi^2} \tilde{W}_{0i} + O(g^4 \ln |t' - t|) \end{aligned}$$

Instantons and Sphalerons

- Vacuum as superposition of vacuum states with different topological winding number, from -infinity up to +infinity

$$|\theta_1, \theta_2\rangle = \sum_m \sum_n e^{i(m\theta_1 + n\theta_2)} |m\rangle_{EW} |n\rangle_{QCD}$$

$$\Delta B = \Delta L = \pm 3n_f$$



- B-L is conserved while B+L is violated
- Fermion levels are shifted in the $|m\rangle$ state relative to the $|m+1\rangle$ state:
 - total „baryon number“ (measured by the gauge invariant current) of each $|m\rangle$ state is zero when we sum over gauge topology and B contributions
 - each $|m\rangle$ state carries zero net electric charge
- Tunneling and vacuum transitions can yield baryon number non-conservation

$$q + q \rightarrow 7\bar{q} + 3\bar{l},$$

Vacuum transition processes

- E-weak instanton tunneling processes strongly suppressed

$$e^{-4\pi \sin^2 \theta_W / \alpha} \sim 10^{-170}$$

- BUT at high temperatures of order the potential barrier (multi TeV) in the early Universe thermal fluctuations can induce vacuum transitions „Sphalerons“ and the suppression factor goes away
- Key equations

$$J_\mu^{\text{con}} = J_\mu - n_f (-K_\mu + k_\mu)$$

$$\Delta Y = \Delta B - n_f m.$$

Vacuum transitions II

- Start from

$$\Delta Y = \Delta B - n_f m.$$

- **Choice of baryon number current essential**
→ yields different and interesting physics

$$\Delta B = n_f \text{ and } \Delta Y = 0$$

- Two solutions (for $m=1$)

OR

$$\Delta Y = -n_f \text{ and } \Delta B = 0.$$


- Essential physics is in zero momentum modes.
 - Vacuum transitions involve zero momentum physics !
 - B+L violation goes for finite-momentum fermions
 - Zero momentum modes (vacuum) restores conservation laws
- The B definition → formation of a zero-momentum „topological condensate“ which accompanies the change in B baryon number
- For the Y charge definition one needs to introduce a zero mode „schizon“ which absorbs B charge in the vacuum → no net condensate

The real world: QCD + E-weak

- E-weak Sphalerons involve only left handed fermions
- Also have QCD Sphalerons plus scalar Higgs couplings → flip the spin/chiralities of the left handed quarks produced by the e-weak sphalerons.
- **Net result is spin independent baryon number violation PLUS spin independent „topological condensate“**
 - » **vector component associated with vector current**
- **(Presumably) still there today with accompanying B violation !**
 - **Phenomenological and cosmological implications ??**

QCD: the Spin Structure of the Proton

- Polarized Deep Inelastic Scattering
 - Measure g_1 spin structure function
 - First moment \rightarrow Quark spin content ~ 0.3
 - Where is the „missing spin“ ?
 - Spawned vast EP program-many exciting ideas being tested: gluon, valence and sea quark polarization ... (COMPASS, RHIC, HERMES)
 - Key result [SDB]:
Transition from current to constituent quarks
 \rightarrow Polarized Condensate ($x=0$)
- Testable through elastic neutrino-proton scattering (measures everything, including $x=0$ contributions)



Conclusions

- Topological condensate: the vacuum has structure
- Spin independent, vector component \rightarrow SSB of Lorentz invariance (?)
- Sign of baryon number violation in sphaleron models
- QCD version of this has important phenomenology in the proton spin and axial U(1) problems